

On 2-Detour Subgraphs of the Hypercube

József Balogh^{1*}, Alexandr Kostochka^{1,2**}

¹ University of Illinois at Urbana-Champaign, IL 61801, USA. e-mail: jobal@math.uiuc.edu

² Institute of Mathematics, Novosibirsk 630090, Russia. e-mail: kostochk@math.uiuc.edu

Abstract. A spanning subgraph H of a graph G is a 2-detour subgraph of G if for each $x, y \in V(G)$, $d_H(x, y) \leq d_G(x, y) + 2$. We prove a conjecture of Erdős, Hamburger, Pippert, and Weakley by showing that for some positive constant c and every n , each 2-detour subgraph of the n -dimensional hypercube Q_n has at least $c \log_2 n \cdot 2^n$ edges.

Key words. Graph, Extremal subgraph, Hypercube, Detour subgraph, Additive spanner.

1. Introduction

Let $d_G(x, y)$ denote the distance between vertices x and y in the graph G . A spanning subgraph H of a graph G is a k -additive spanner if for each pair (u, v) of vertices of G , $d_H(u, v) \leq d_G(u, v) + k$. Studying k -additive spanners was motivated by a number of problems in communication networks, broadcasting, routing, etc., see [7, 9, 10]. Additive spanners were studied in [1–6, 8]. Sometimes, k -additive spanners of the n -dimensional hypercube Q_n are also called k -detour subgraphs.

Let $f_k(n)$ denote the minimum number of the edges of a k -detour subgraph of Q_n . Since Q_n is a bipartite graph, it is enough to consider $f_k(n)$ only for even k . Erdős, Hamburger, Pippert, and Weakley [3] studied 2-detour subgraphs of Q_n . They constructed a 2-detour subgraph of Q_n with at most $3/(2\sqrt{2})\sqrt{n}2^n$ edges. Since each k -detour subgraph of Q_n is connected, it has at least $2^n - 1$ edges. The best known lower bound on $f_2(n)$ is due to Hamburger, Kostochka and Sidorenko [4], who proved that

$$f_2(n) \geq (3.000013 - o(1)) \cdot 2^n. \quad (1)$$

Arizumi, Hamburger and Kostochka [1] proved that $f_4(n) < (3 + o(1)) \cdot 2^n$ for large n , that is, that the average degree of some 4-detour subgraphs of Q_n is less than 7 for large n . In contrast, it was conjectured in [3] that the function $f_2(n) \cdot 2^{-n}$ is unbounded. The main result of our note is a proof of the conjecture.

* Research supported in part by NSF grants DMS-0302804, DMS-0603769 and DMS-0600303, UIUC Campus Reseach Board #06139 and #07048, and OTKA 049398.

** Research supported in part by NSF grants DMS-0400498 and DMS-0650784, and grant 06-01-00694 of the Russian Foundation for Basic Research.

Theorem 1. *If $H = H_n$ is a 2-detour subgraph of the n -dimensional hypercube Q_n , then*

$$2e(H) > 10^{-6} 2^n \log_2 n. \tag{2}$$

The idea of the proof is the following: If H is a 2-detour subgraph of Q_n with $10^{-6} 2^n \log_2 n$ edges, then we claim several properties of H . First, we claim that ‘most’ vertices of H have a ‘low’ degree. Then we show that for some special m between ‘most’ pairs of low vertices at distance m in Q_n the shortest path in H has length $m + 2$, additionally the first and the last edges in these paths are ‘parallel’. After claiming these properties, a simple counting argument provides the result. A difficulty arises from that we do not know the precise value of this special m , we could only prove its existence.

We introduce the necessary notation in the next section and prove a series of preparatory statements in Section 3. Theorem 1 will be proved in Section 4.

2. Notation

Let $\ell = \lfloor \log_2 n \rfloor$. Let $H = H_n$ be a 2-detour subgraph of Q_n , with

$$e(H) = t \cdot 2^n \tag{3}$$

for some t . By the definition of a 2-detour subgraph, H has no isolated vertices. By $d(v)$ we denote the degree of v in H , but by $d(u, v)$ we denote the distance between u and v in Q_n . For $i = 0, \dots, \ell$, let

$$A_i := \{x \in V(H) : 2^i \leq d(x) < 2^{i+1}\}, \quad \tilde{A}_i := \bigcup_{j=i}^{\ell} A_j, \tag{4}$$

$$a_i := |A_i| \quad \text{and} \quad \tilde{a}_i := |\tilde{A}_i|. \tag{5}$$

By the definition of a_i ,

$$\sum_{i=0}^{\ell} a_i \cdot 2^i \leq 2e(H) < \sum_{i=0}^{\ell} a_i \cdot 2^{i+1}. \tag{6}$$

We say that a vertex x is *low* if $d(x) \leq 2000t$. The set of low vertices is

$$L := \{x : d(x) \leq 2000t\}. \tag{7}$$

It is convenient to consider the vertices in a vector form, where they are $\{0, 1\}$ -vectors of length n . The coordinates of a vertex sometimes are referred to as *directions*.

For each two vertices $u, v \in V(H)$, we define

$$\text{co}(u, v) := \{\text{the set of coordinates where } u \text{ and } v \text{ differ}\}. \tag{8}$$

Note that $d(u, v) = |\text{co}(u, v)|$ and that each (u, v) -path uses the edges of each direction in $\text{co}(u, v)$ an odd number of times. For a vertex u and a positive integer r , let

$$\text{Dir}(u, r) := \{\text{co}(u, v) : v \in N(u)\} \cup \{\text{co}(v, w) : v \in N(u), d(v) < 2^r, w \in N(v)\}. \tag{9}$$

In other words, $\text{Dir}(u, r)$ is the union of the directions (in H) from u toward its neighbors, and the directions from not-very-high degree neighbors of u towards their neighbors. For a vertex u and positive integers m and r , we define the set $S(u, m, r)$ of vertices at distance m from u such that the shortest paths from u to them do not use any direction from $\text{Dir}(u, r)$: Let

$$B(u, m) := \{v : d(u, v) = m\} \tag{10}$$

be the sphere of radius m about vertex u and

$$S(u, m, r) := \{x \in B(u, m) : \text{co}(u, x) \cap \text{Dir}(u, r) = \emptyset\}. \tag{11}$$

Finally, let $\tilde{S}(u, m, r) := \{v \in S(u, m, r) : u \in S(v, m, r)\}$.

3. Preliminaries

Proposition 1.

$$|L| > \frac{999}{1000} \cdot 2^n.$$

Proof. The chain of inequalities

$$2t \cdot 2^n = \sum_{x \in V(Q_n)} d(x) > \sum_{x \in V(Q_n) \setminus L} d(x) > 2000t(2^n - |L|)$$

implies our claim. □

The next observation follows from the definitions of L and Dir .

Proposition 2. *Let $u \in L$ and $r \leq \ell$ be a positive integer. Then*

$$|\text{Dir}(u, r)| < 2000 \cdot t \cdot 2^r.$$

□

For each $u \in L$, let

$$T(u) := \{r \in \{0, 1, 2, \dots, \ell\} : |\text{Dir}(u, r)| \leq 400000 \cdot t \cdot 2^r / \ell\}. \tag{12}$$

Proposition 3. *For each $u \in L$, $|T(u)| \geq 0.99(\ell + 1)$.*

Proof. Consider $\phi(u) := \sum_{r=0}^{\ell} |\text{Dir}(u, r)|2^{-r}$. A neighbor $v \in A_i$ of u contributes to the summand $|\text{Dir}(u, r)|2^{-r}$ the amount of $d(v)2^{-r} < 2^{i+1-r}$ if $i \leq r - 1$, and contributes 0 otherwise. Hence, in total v contributes less than 2 to $\phi(u)$. Therefore,

$$\phi(u) = \sum_{r=0}^{\ell} |\text{Dir}(u, r)|2^{-r} < 2d(u) \leq 4000t. \tag{13}$$

In order (13) to hold, fewer than 0.01ℓ summands can exceed $400000t/\ell$. □

For each $r \in \{0, 1, 2, \dots, \ell\}$, let

$$m(r) := \left\lfloor \min \left\{ 0.1n, 10^{-8} \frac{\ell \cdot n}{t} 2^{-r} \right\} \right\rfloor. \tag{14}$$

Proposition 3 will be used to prove the following two claims.

Proposition 4. *Let $u \in L$. Then for each $r \in T(u)$,*

$$|S(u, m(r), r)| \geq 0.99 \binom{n}{m(r)}. \tag{15}$$

Proof. Let $u \in L, r \in T(u)$ and $m = m(r)$. By the definitions of $T(u)$ and $S(u, m(r), r)$,

$$\begin{aligned} \frac{|S(u, m, r)|}{\binom{n}{m}} &= \frac{\binom{n - |\text{Dir}(u, r)|}{m}}{\binom{n}{m}} \geq \frac{\binom{n - 4 \cdot 10^5 t \cdot 2^r / \ell}{m}}{\binom{n}{m}} \geq \left(\frac{n - m - 4 \cdot 10^5 t \cdot 2^r / \ell}{n - m} \right)^m \\ &\geq 1 - m \frac{4 \cdot 10^5 t \cdot 2^r}{\ell(n - m)}. \end{aligned}$$

By the definition of $m = m(r)$, we have

$$m \frac{4 \cdot 10^5 t \cdot 2^r}{\ell(n - m)} \leq \frac{\ell \cdot n \cdot 4 \cdot 10^5 \cdot t \cdot 2^r}{10^8 t \ell(n - m) 2^r} = \frac{0.004n}{n - m} \leq 0.01.$$

This proves the proposition. □

Proposition 5. *For at least $0.5 \cdot 2^n$ vertices $u \in L$, there are at least $0.8(\ell + 1)$ values of $r \in \{0, 1, 2, \dots, \ell\}$ such that*

$$|\{v \in B(u, m(r)) : u \in S(v, m(r), r)\}| > 0.75 \binom{n}{m(r)}. \tag{16}$$

Proof. Propositions 3 and 4 imply that

$$\sum_{v \in L} \sum_{r=0}^{\ell} \frac{|S(v, m(r), r)|}{\binom{n}{m(r)}} \geq (0.99)^2 (\ell + 1) |L| > 0.98 (\ell + 1) |L|.$$

By Proposition 1, the last expression is greater than $0.979(\ell + 1)2^n$. But this sum is less than

$$\sum_{u \in V(H)} \sum_{r=0}^{\ell} \frac{|\{v \in B(u, m(r)) : u \in S(v, m(r), r)\}|}{\binom{n}{m(r)}}.$$

It follows that for at least $0.51 \cdot 2^n$ vertices $u \in V(H)$,

$$\sum_{r=0}^{\ell} \frac{|\{v \in B(u, m(r)) : u \in S(v, m(r), r)\}|}{\binom{n}{m(r)}} > 0.95(\ell + 1). \tag{17}$$

Hence, by Proposition 1, (17) holds for at least $0.5 \cdot 2^n$ vertices $u \in L$. And (17) cannot hold for a vertex $u \in L$ if for more than $0.2(\ell + 1)$ values of $r \in \{0, 1, 2, \dots, \ell\}$, (16) fails. \square

By (6),

$$\tilde{a}_r = |\cup_{j=r}^{\ell} A_j| \leq 2e(H) \cdot 2^{-r} = t \cdot 2^{n-r+1}. \tag{18}$$

Let

$$R := \{r \in \{0, 1, 2, \dots, \ell\} : \tilde{a}_r < 20t \cdot 2^{n-r}/\ell\}. \tag{19}$$

Proposition 6. $|R| \geq 4(\ell + 1)/5$.

Proof. Let $S := \sum_{r=0}^{\ell} 2^r \tilde{a}_r$. By definition of \tilde{a}_r and by (6),

$$S = \sum_{r=0}^{\ell} 2^r \sum_{j=r}^{\ell} a_j = \sum_{j=0}^{\ell} a_j \sum_{r=0}^j 2^r < \sum_{j=0}^{\ell} a_j 2^{j+1} \leq 2 \sum_{u \in V(H)} d(u) = 4t \cdot 2^n.$$

In order S to be less than $4t \cdot 2^n$, fewer than $\ell/5$ summands can be greater or equal to $20t \cdot 2^n/\ell$. \square

Proposition 7. For each $r \in R$ and for any given $1 < m < n$,

$$\left| \left\{ x : |B(x, m) \cap \tilde{A}_r| \leq 20000 \frac{t}{\ell} 2^{-r} \binom{n}{m} \right\} \right| > \frac{999}{1000} \cdot 2^n. \tag{20}$$

Proof. Observe that for each r and m ,

$$\sum_{x \in V(H)} |B(x, m) \cap \tilde{A}_r| = \tilde{a}_r \binom{n}{m}. \tag{21}$$

Hence, if $r \in R$, then the number of summands on the left-hand side of (21) exceeding $20000 \frac{t}{\ell} 2^{-r} \binom{n}{m}$ is less than $10^{-3} 2^n$. \square

Proposition 7 immediately yields the following.

Proposition 8. Let $m(r)$ be defined by (14). For each $r \in R$, the number of vertices $x \in V(H)$ such that

$$|B(x, m(r) + 1) \cap \tilde{A}_r| \leq 20000 \frac{t}{\ell} 2^{-r} \binom{n}{m(r) + 1} \tag{22}$$

is at least $0.999 \cdot 2^n$. \square

We extend it as follows.

Proposition 9. *At least $0.99 \cdot 2^n$ vertices $x \in V(H)$ possess the following property: For at least $3(\ell + 1)/5$ values of $r \in \{0, 1, 2, \dots, \ell\}$,*

$$|B(x, m(r) + 1) \cap \tilde{A}_r| \leq 20000 \frac{t}{\ell} 2^{-r} \binom{n}{m(r) + 1}.$$

Proof. For each $r \in R$, let $X(r)$ be the set of vertices of H for which (22) does not hold. By Proposition 8, $|X(r)| \leq 0.001 \cdot 2^n$ for each $r \in R$. Hence $\sum_{r \in R} |X(r)| \leq 0.001 \cdot 2^n |R|$ and the number of vertices $v \in V(H)$ that belong to at least $(\ell + 1)/5$ sets $X(r)$ is at most

$$\frac{0.001 \cdot 2^n |R|}{0.2(\ell + 1)} \leq 0.01 \cdot 2^n.$$

Since $|R| \geq 4(\ell + 1)/5$, this proves the proposition. □

The next easy observation is one of our key tools.

Proposition 10. *Let vertices $u, v \in L$ be such that $v \in \tilde{S}(u, m, r)$ for some integers $m \geq 4$ and $r \geq 1$. Let P be a (u, v) -path in H of length at most $m + 2$, with the first edge ux and the last edge yv . Then the following statements hold:*

- (i) *The length of P is exactly $m + 2$;*
- (ii) *$x, y \in \tilde{A}_r$;*
- (iii) *$\text{co}(u, x) = \text{co}(v, y)$;*
- (iv) *$y \in B(u, m(r) + 1) \cap \tilde{A}_r$, and $\text{co}(y, v) \in \text{Dir}(u, r)$;*
- (v) *$\text{co}(y, u) = \text{co}(v, u) \cup \text{co}(y, v)$.*

Proof. By the definition of $\tilde{S}(u, m, r)$, $\text{co}(u, x) \not\subseteq \text{co}(u, v)$ and $\text{co}(v, y) \not\subseteq \text{co}(u, v)$. This implies that both edges, ux and vy , are additional to the shortest (u, v) -path in Q_n , implying (i). Furthermore, if $\text{co}(u, x) \neq \text{co}(v, y)$, then extra edges are needed in P for both directions, which makes the length of P at least $m + 4$, a contradiction, yielding (iii).

If $x \notin \tilde{A}_r$ and the second edge of P is xz , then $\text{co}(x, z) \in \text{Dir}(u, r)$, and hence by (11), the edge xz is extra to make P longer than $m + 2$. Thus, $x \in \tilde{A}_r$. Similar argument proves $y \in \tilde{A}_r$, implying (ii). Part (ii) already gives $y \in \tilde{A}_r$. As $\text{co}(y, v) \not\subseteq \text{co}(u, v)$ we have that $d(u, y) = m(r) + 1$, yielding instantly (iv) and (v). □

4. Proof of the Main Result

Choose a vertex $u \in L$ for which the statements of Propositions 9 and 5 hold. Then by Propositions 3, 4, 9, and 5, there are at least $(1 - 0.01 - 0.4 - 0.2)(\ell + 1) = 0.39(\ell + 1)$ values of $r \in \{0, 1, 2, \dots, \ell\}$ such that

$$|B(u, m(r) + 1) \cap \tilde{A}_r| \leq 20000 \frac{t}{\ell} 2^{-r} \binom{n}{m(r) + 1} \tag{23}$$

and (recalling that $\tilde{S}(u, m, r) = \{v \in S(u, m, r) : u \in S(v, m, r)\}$ and combining (15) and (16))

$$|\tilde{S}(u, m(r), r)| > (0.75 - 0.01) \binom{n}{m(r)}. \tag{24}$$

Choose some $r \in \{0, 1, 2, \dots, \ell\}$ satisfying (23) and (24) with $r \geq 0.1\ell$. If n (and hence ℓ) is large enough, then $m(r) = \lfloor 10^{-8} \frac{\ell \cdot n}{t} 2^{-r} \rfloor$.

Let $v \in \tilde{S}(u, m(r), r)$. By the definition of H , there is a (u, v) -path P_{uv} of length at most $m(r) + 2$. By Proposition 10 (iv), the neighbor $y = y(v)$ of v on the path P_{uv} is in $B(u, m(r) + 1) \cap \tilde{A}_r$, and $\text{co}(y, v) \in \text{Dir}(u, r)$. By the definition of $\tilde{S}(u, m(r), r)$ we have that $\text{co}(v, u) \cap \text{Dir}(u, r) = \emptyset$. Proposition 10 (v) states that $\text{co}(y, u) = \text{co}(v, u) \cup \text{co}(y, v)$, in particular that $\text{co}(y, u) \cap \text{Dir}(u, r) = \text{co}(y, v)$. Therefore, given u and y , the direction $\text{co}(y, v)$ is determined. So the only vertex in $\tilde{S}(u, m(r), r)$ that can be reached from u by a path of length $m(r) + 2$ passing vertex y is v . In other words, the number of choices for v is not larger than for y , i.e.,

$$|\tilde{S}(u, m(r), r)| \leq |B(u, m(r) + 1) \cap \tilde{A}_r|.$$

Using (23) and (24), we get

$$0.74 \binom{n}{m(r)} < 20000 \frac{t}{\ell} 2^{-r} \binom{n}{m(r) + 1}.$$

Hence,

$$\frac{m(r) + 1}{n - m(r)} < 30000 \frac{t}{\ell} 2^{-r}.$$

Plugging in the value $m(r) = \lfloor 10^{-8} \frac{\ell \cdot n}{t} 2^{-r} \rfloor$ from the previous paragraph, we have

$$\frac{10^{-8} \frac{\ell \cdot n}{t} 2^{-r}}{n} < \frac{m(r) + 1}{n - m(r)} < 30000 \frac{t}{\ell} 2^{-r}.$$

This yields

$$t > \frac{1}{\sqrt{3}} 10^{-6} \ell = \frac{1}{\sqrt{3}} 10^{-6} \lfloor \log_2 n \rfloor > 0.5 \cdot 10^{-6} \log_2 n,$$

which proves our theorem, since $t = 2^{-n} e(H)$. □

Acknowledgements. We thank the referee for careful reading of the first version of the manuscript.

References

1. Arizumi, N., Hamburger, P., Kostochka, A.V.: On k -detour subgraphs of hypercubes. *J. Graph Theory* **57**, 55–64 (2008)
2. Chepoi, V., Dragan, F., Yan, C.: Additive spanners for k -chordal graphs, in *Algorithms and Complexity*, 96–107, Springer LNCS **2653**, 2003
3. Erdős, P., Hamburger, P., Pippert, R.E., Weakley, W.D.: Hypercube subgraphs with minimal detours. *J. Graph Theory* **23**, 119–128 (1996)
4. Hamburger, P., Kostochka, A.V., Sidorenko, A.: Hypercube subgraphs with local detours. *J. Graph Theory* **30**, 101–111 (1999)
5. Kratsch, D., Le, H., Müller, H., Prisner, E., Wagner, D.: Additive tree spanners. *SIAM J. Discrete Math.* **17**(2), 332–340 (2004)
6. Liestman, A.L., Shermer, T.C.: Additive graph spanners. *Networks* **23**(4), 343–363 (1993)
7. Peleg, D., Schäffer, A.A.: Graph spanners. *J. Graph Theory* **13**(1), 99–116 (1989)
8. Peleg, D., Ullman, J.D.: An optimal synchronizer for the hypercube. in *Proceedings of the Sixth Annual ACM Symposium on Principles of Distributed Computing*, 77–85 (1987)
9. Soares, J.: Graph spanners: A survey. *Congressus Numerantium* **89**, 225–238 (1992)
10. Venkatesan, G., Rotics, U., Madanlal, M.S., Makowsky, J.A., Pandu Rangan, C.: Restrictions of minimum spanner problems. *Inform. and Comput.* **136**(2), 143–164 (1997)

Received: May 30, 2006

Final Version received: May 6, 2008