

# Graphs with maximum degree 5 are acyclically 7-colorable\*

Alexandr V. Kostochka<sup>†</sup>

*Department of Mathematics, University of Illinois, Urbana, IL 61801, USA*

*Sobolev Institute of Mathematics, Novosibirsk, 630090, Russia*

Christopher Stocker

*Department of Mathematics, University of Illinois, Urbana, IL 61801, USA*

Received 12 May 2010, accepted 21 January 2011, published online xx March 2011

---

## Abstract

An acyclic coloring is a proper coloring with the additional property that the union of any two color classes induces a forest. We show that every graph with maximum degree at most 5 has an acyclic 7-coloring. We also show that every graph with maximum degree at most  $r$  has an acyclic  $(1 + \lfloor \frac{(r+1)^2}{4} \rfloor)$ -coloring.

*Keywords:* Acyclic coloring, maximum degree.

*Math. Subj. Class.:* 05C15, 05C35

---

## 1 Introduction

A proper coloring of the vertices of a graph  $G = (V, E)$  is an assignment of colors to the vertices of the graph such that no two adjacent vertices receive the same color. A proper coloring of a graph  $G$  is *acyclic* if the union of any two color classes induces a forest. The *acyclic chromatic number*,  $a(G)$ , is the smallest integer  $k$  such that  $G$  is acyclically  $k$ -colorable. The notion of acyclic coloring was introduced in 1973 by Grünbaum [8] and turned out to be interesting and closely connected to a number of other notions in graph coloring. Several researchers felt the beauty of the subject and started working on problems and conjectures posed by Grünbaum. Michael Albertson was among the enthusiasts and wrote in total four papers on the topic [1, 2, 3, 4].

---

\*Dedicated to the memory of Michael Albertson

<sup>†</sup>Supported in part by NSF grants DMS-0650784 and DMS-0965587 and grant 09-01-00244-a of the Russian Foundation for Basic Research.

*E-mail addresses:* kostochk@math.uiuc.edu (Alexandr V. Kostochka), stocker2@illinois.edu (Christopher Stocker)

In particular, Grünbaum studied  $a(r)$  – the maximum value of the acyclic chromatic number over all graphs  $G$  with maximum degree at most  $r$ . He conjectured that for every  $r$ ,  $a(r) = r + 1$  and proved that his conjecture holds for  $r \leq 3$ . In 1979, Burstein [6] proved the conjecture for  $r = 4$ . This result was proved independently by Kostochka [10]. It was also proved in [10] that for every  $k \geq 3$ , the problem of deciding whether a graph is acyclically  $k$ -colorable is NP-complete. It turned out that for large  $r$ , Grünbaum's conjecture is incorrect in a strong sense. Albertson and Berman mentioned in [1] that Erdős proved that  $a(r) = \Omega(r^{4/3-\epsilon})$  and conjectured that  $a(r) = o(r^2)$ . Alon, McDiarmid and Reed [5] sharpened Erdős' lower bound to  $a(r) \geq cr^{4/3}/(\log r)^{1/3}$  and proved that

$$a(r) \leq 50r^{4/3}. \quad (1.1)$$

This established almost the order of the magnitude of  $a(r)$  for large  $r$ . Recently, the problem of estimating  $a(r)$  for small  $r$  was considered again.

Fertin and Raspaud [7] showed among other results that  $a(5) \leq 9$  and gave a linear-time algorithm for acyclic 9-coloring of any graph with maximum degree 5. Furthermore, for every fixed  $r \geq 3$ , they gave a fast algorithm that uses at most  $r(r-1)/2$  colors for acyclic coloring of any graph with maximum degree  $r$ . Of course, for large  $r$  this is much worse than the upper bound (1.1), but for  $r < 1000$ , it is better. Hocquard and Montassier [9] showed that every 5-connected graph  $G$  with  $\Delta(G) = 5$  has an acyclic 8-coloring. Kothapalli, Varagani, Venkaiah, and Yadav [12] showed that  $a(5) \leq 8$ . Kothapalli, Satish, and Venkaiah [11] proved that every graph with maximum degree  $r$  is acyclically colorable with at most  $1 + r(3r+4)/8$  colors. This is better than the bound  $r(r-1)/2$  in [7] for  $r \geq 8$ . The main result of this paper is

**Theorem 1.1.** *Every graph with maximum degree 5 has an acyclic 7-coloring, i.e.,  $a(5) \leq 7$ .*

We do not know whether  $a(5)$  is 7 or 6, and do not have a strong opinion about it.

Our proof is different from that in [7, 9, 12] and heavily uses the ideas of Burstein [6]: he started from an uncolored graph  $G$  with maximum degree 4 and colored step by step more and more vertices (with some recolorings) so that each of partial acyclic 5-colorings of  $G$  had additional good properties that enabled him to extend the coloring further. The proof yields a linear-time algorithm for acyclic coloring with at most 7 colors of any graph with maximum degree 5. Using this approach we also show that for every fixed  $r \geq 6$ , there exists a linear-time algorithm giving an acyclic coloring of any graph with maximum degree  $r$  with at most  $1 + \lfloor \frac{(r+1)^2}{4} \rfloor$  colors. This is better than the bounds in [7] and [11] cited above for every  $r \geq 6$ .

In the next section we introduce notation, prove two small lemmas and state the main lemma. In Section 3 we prove Theorem 1.1 modulo the main lemma. In Section 4 we derive linear-time algorithms for acyclic coloring of graphs with bounded maximum degree. In the last section we give the proof of the main lemma.

## 2 Preliminaries

Let  $G$  be a graph. A *partial coloring* of  $G$  is a coloring of some subset of the vertices of  $G$ . A *partial acyclic coloring* is then a proper partial coloring of  $G$  containing no bicolored cycles.

- 58 Given a partial coloring  $f$  of  $G$ , a vertex  $v$  is  
 59 (a) *rainbow* if all colored neighbors of  $v$  have distinct colors;  
 60 (b) *almost rainbow* if there is a color  $c$  such that exactly two neighbors of  $v$  are colored  
 61 with  $c$  and all other colored neighbors of  $v$  have distinct colors;  
 62 (c) *admissible* if it is either rainbow or almost rainbow;  
 63 (d) *defective* if  $v$  is an uncolored almost rainbow vertex such that at least one of the two of  
 64 its neighbors receiving the same color is admissible.

65 A partial acyclic coloring  $f$  of a graph  $G$  is *rainbow* if  $f$  is a partial acyclic coloring of  
 66  $G$  such that every uncolored vertex is rainbow.

67 A partial acyclic coloring  $f$  of a graph  $G$  is *admissible* if either  $f$  is rainbow or one  
 68 vertex is defective and all other uncolored vertices are rainbow. In these terms, a coloring  
 69 is rainbow if it is admissible and has no defective vertices. Note that both, rainbow and  
 70 admissible colorings are partial acyclic colorings where additional restrictions are put only  
 71 on uncolored vertices. The advantage of using admissible colorings is that they provide a  
 72 stronger induction condition that places additional restrictions only on coloring of neigh-  
 73 bors of uncolored vertices. So, the fewer uncolored vertices remains, the weaker are these  
 74 additional restrictions.

75 All colorings in this section will be from the set  $\{1, 2, \dots, 7\}$ .

76 **Lemma 2.1.** *Let  $v$  be a vertex of degree 4 in a graph  $G$  with  $\Delta(G) \leq 5$ . Let  $f$  be an*  
 77 *admissible (respectively, rainbow) coloring in which  $v$  is colored with color  $c_1$ , each of the*  
 78 *neighbors of  $v$  is colored, and exactly 3 colors appear on the neighbors of  $v$ . If at least*  
 79 *one of the two neighbors of  $v$  receiving the same color and one of the other two neighbors*  
 80 *of  $v$  each have a second (i.e., distinct from  $v$ ) neighbor with color  $c_1$ , then we can recolor*  
 81  *$v$  and at most one of its neighbors so that the coloring remains admissible (respectively,*  
 82 *rainbow). In particular, the new partial acyclic coloring has no new defective vertices.*  
 83 *Moreover, if we need to recolor a vertex other than  $v$ , then we may choose a vertex with 5*  
 84 *colored neighbors and recolor it with a color incident to  $v$  in  $f$ .*

85 *Proof.* Let  $N(v) = \{z_1, z_2, z_3, z_4\}$ ,  $f(z_1) = f(z_2) = c_2$ ,  $f(z_3) = c_3$ ,  $f(z_4) = c_4$ . Let  $z_2$   
 86 and  $z_3$  be the neighbors of  $v$  with colors  $c_2$ , and  $c_3$  that are also adjacent to another vertex  
 87 of color  $c_1$ . We may assume that  $z_2$  is adjacent to a vertex of color  $c_5$ , since otherwise  
 88 when we recolor  $v$  with  $c_5$ , no bicolored cycles appear and the coloring remains admissible  
 89 (respectively, rainbow). Similarly, we may assume that  $z_2$  is adjacent to vertices of colors  
 90  $c_6$  and  $c_7$ . Then we may recolor  $z_2$  with  $c_3$  and repeat the above argument to get that  $z_3$  also  
 91 is adjacent to vertices with colors  $c_5$ ,  $c_6$ , and  $c_7$ . In this case, we may change the original  
 92 coloring by recoloring  $z_3$  with  $c_2$  and  $v$  with  $c_3$ . So, in this case only  $v$  and  $z_3$  change  
 93 colors. Note that either only  $v$  changes its color, or  $z_2$  receives color  $c_3$ , or  $z_3$  receives  
 94 color  $c_2$ . □

95 For partial colorings  $f$  and  $f'$  of a graph  $G$ , we say that  $f'$  is *larger than*  $f$  if it colors  
 96 more vertices.

97 **Lemma 2.2.** *Let  $v$  be a vertex of degree 4 in a graph  $G$  with  $\Delta(G) \leq 5$ . Let  $f$  be a rainbow*  
 98 *coloring in which  $v$  is colored with color  $c_1$ , the neighbors  $z_1$ ,  $z_2$ , and  $z_3$  of  $v$  receive the*  
 99 *distinct colors  $c_2$ ,  $c_3$ , and  $c_4$ , the neighbor  $z_4$  of  $v$  is an uncolored rainbow vertex. Then*  
 100 *either  $G$  has a rainbow coloring  $f_1$  that colors the same vertices and differs from  $f$  only*  
 101 *at  $v$ , or  $G$  has a rainbow coloring  $f'$  larger than  $f$ . Moreover, if the former does not hold,*

102 then  $z_4$  has degree 5 and exactly one uncolored neighbor, say  $z_{4,4}$ , and we can choose the  
103 larger coloring  $f'$  so that all the following are true:

- 104 1. Every vertex colored in  $f$  is still colored.
- 105 2. Vertex  $z_4$  is colored.
- 106 3. The only uncolored vertex apart from  $z_4$  that may get colored is  $z_{4,4}$ , and it does only  
107 if it has neighbors of colors  $c_1, c_2, c_3$ , and  $c_4$ .
- 108 4. Apart from  $v$ , only one vertex  $w$  may change its color, and if it does, then (a)  $w$  is a  
109 neighbor of  $z_4$ , (b)  $w$  has four colored neighbors, (c) it changes a color in  $\{c_5, c_6, c_7\}$   
110 to another color in  $\{c_5, c_6, c_7\}$ , and (d)  $z_4$  gets the former color of  $w$ . In particular,  
111  $v$  is admissible in  $f'$ .

112 *Proof.* Let  $v, z_1, z_2, z_3$ , and  $v_4$  be as in the hypothesis. We may assume that  $z_4$  is adjacent  
113 to a vertex  $z_{4,1}$  of color  $c_5$ : otherwise, since  $v_4$  is rainbow, when we recolor  $v$  with  $c_5$ ,  
114 the new coloring will be rainbow. Similarly, we may assume that  $z_4$  is adjacent to vertices  
115  $z_{4,2}$ , and  $z_{4,3}$  of colors  $c_6$  and  $c_7$ . If  $z_4$  has no other neighbors, then we can recolor  $v$   
116 with  $c_5$  and color  $z_4$  with  $c_1$ . So, assume that  $z_4$  has the fifth neighbor,  $z_{4,4}$ . If  $z_{4,4}$  is  
117 colored, then  $f(z_{4,4}) \in \{c_2, c_3, c_4\}$ , since  $z_4$  is rainbow. In this case, we let  $f'(z_4) = c_1$   
118 and  $f'(v) = c_5$ . So, we may assume that  $z_{4,4}$  is not colored. If  $z_{4,4}$  has no neighbor of  
119 color  $c_2$ , then coloring  $z_4$  with  $c_2$  leaves the coloring rainbow and makes it larger than  $f$ .  
120 Thus, we may assume that  $z_{4,4}$  has a neighbor of color  $c_2$  and similarly neighbors of colors  
121  $c_3$  and  $c_4$ . If  $z_{4,4}$  has no neighbor of color  $c_1$ , then we let  $f'(z_4) = c_1$  and  $f'(v) = c_5$ . So,  
122 let  $z_{4,4}$  have such a neighbor.

123 If  $z_{4,1}$  has no neighbor of color  $c_2$ , then by coloring  $z_4$  with  $c_2$  and  $z_{4,4}$  with  $c_5$ , we  
124 get a rainbow coloring larger than  $f$ . So, we may assume (by symmetry) that  $z_{4,1}$  has  
125 neighbors of colors  $c_2, c_3, c_4$ . If  $z_{4,1}$  has no neighbor of color  $c_1$ , then we let  $f'(z_4) = c_1$ ,  
126  $f'(z_{4,4}) = c_5$ , and  $f'(v) = c_6$ . Finally, if  $z_{4,1}$  also has a neighbor of color  $c_1$ , then we let  
127  $f'(z_{4,1}) = c_6$  and  $f'(z_4) = c_5$ .  $\square$

128 The next lemma is our main lemma. We will use it in the next section and prove in  
129 Section 5.

130 **Lemma 2.3.** *Let  $f$  be an admissible partial coloring of a 5-regular graph  $G$ . Then  $G$  has  
131 a rainbow coloring  $f'$  that colors at least as many vertices as  $f$ .*

### 132 3 Proof the the Theorem

133 For convenience, we restate Theorem 1.1.

134 **Theorem.** Every graph with maximum degree 5 has an acyclic 7-coloring.

135 *Proof.* Let  $G$  be such a graph. If  $G$  is not 5-regular, form  $G'$  from two disjoint copies  
136 of  $G$  by adding for each  $v \in V(G)$  of degree less than 5 an edge between the copies of  
137  $v$ . Repeating this process at most five times gives a 5-regular graph  $G^*$  containing  $G$  as a  
138 subgraph. Since an acyclic 7-coloring of  $G^*$  yields an acyclic 7-coloring of its subgraph  
139  $G$ , we may assume that  $G$  is 5-regular.

140 Let  $f$  be an admissible coloring of  $G$  from the set  $\{1, 2, \dots, 7\}$  with the most colored  
141 vertices. By Lemma 2.3, we may assume that  $f$  is rainbow.

Let  $H$  be the subgraph of  $G$  induced by the vertices left uncolored by  $f$ . Let  $x$  be a vertex of minimum degree in  $H$ . We consider several cases according to the degree  $d_H(x)$ .

**Case 1:**  $d_H(x) = 0$ . Since  $f$  is rainbow, any color in  $\{1, 2, \dots, 7\} - f(N_G(x))$  can be used to color  $x$  contradicting the maximality of  $f$ .

**Case 2:**  $d_H(x) = 1$ . Since  $f$  is rainbow, we may assume that  $x$  is adjacent to vertices of colors 1, 2, 3, and 4. Let  $y$  be the uncolored neighbor of  $x$ . Since  $y$  is rainbow, coloring  $x$  with 5 gives either a rainbow coloring or an admissible coloring with the defective vertex  $y$  having the admissible neighbor  $x$ , a contradiction to the maximality of  $f$ .

**Case 3:**  $d_H(x) = 2$ . We may assume that  $x$  is adjacent to vertices with colors 1, 2, 3, and two uncolored vertices  $y_1$  and  $y_2$ . Since in our case  $y_1$  is adjacent to at most 3 colored vertices, some color  $c \in \{4, 5, 6, 7\}$  does not appear on the neighbors of  $y_1$ . Coloring  $x$  with  $c$  then yields either a rainbow coloring, or an admissible coloring with defective vertex  $y_2$  and its admissible neighbor  $x$ , a contradiction to the maximality of  $f$ .

**Case 4:**  $d_H(x) = 3$ . We may assume that  $x$  is adjacent to vertices of colors 1 and 2. By the choice of  $x$ , each uncolored vertex of  $G$  has at most 2 colored neighbors. Since the three uncolored neighbors of  $x$  have at most 6 colored neighbors in total, some color  $c \in \{3, 4, 5, 6, 7\}$  is present at most once among these 6 neighbors. Then coloring  $x$  with  $c$  again yields an admissible coloring, a contradiction to the maximality of  $f$ .

**Case 5:**  $d_H(x) \geq 4$ . Since each vertex of  $G$  has at most one colored neighbor, at most 5 colors are used in the second neighborhood of  $x$ . Hence  $x$  may be colored to give a rainbow coloring with more colored vertices.

We conclude that  $H$  is empty and that  $f$  is an acyclic 5-coloring of  $G$ .  $\square$

## 4 Algorithms

**Theorem 4.1.** *There exists a linear time algorithm for finding an acyclic 7-coloring of a graph with maximum degree 5.*

*Proof.* The proof of the Theorem 1.1, along with Lemmas 2.1–2.3 gives an algorithm. In order to control the efficiency of the algorithm we make the following modification: whenever the proof checks whether a vertex  $v$  is in a two-colored cycle, we check only for such a cycle of length at most 12, and if we do not find such a short cycle, then check whether two bicolored paths of length 6 leave  $v$ . This is enough, since the existence of such paths already makes the proofs of Theorem 1.1 and all the lemmas work. So, we need only to consider a bounded (at most  $5^6$ ) number of vertices around our vertex. It then suffices to compute the running time of this algorithm. Let  $n$  be the number of vertices in  $G$ . The process of creating a 5-regular graph takes  $O(n)$  time since we apply this process at most 5 times, each time on at most  $2^5 n$  vertices, each of degree at most 5. We may now assume that  $G$  is a 5-regular graph. We then create and maintain 6 databases  $D_j$ ,  $j = 0, 1, \dots, 5$  (say doubly linked lists), each for the set of vertices with degree  $j$  in the current  $H$ . At the beginning, all vertices are in  $D_5$ , and it is possible to update the databases in a constant amount of time each time a vertex gains or loses a colored neighbor. Since there are at most  $2^5 n$  possible searches for a vertex with the minimum number of uncolored neighbors, all the searches and updates will take  $O(n)$  time. Note that the processes of Lemma 2.1 and Lemma 2.2 also take a constant amount of time to complete. Observe that each of the cases in Lemma 2.3 either finds a rainbow coloring, or finds an admissible coloring with more colored vertices, or reduces to a previous case in an amount of time bounded by a constant. Also when Lemma 2.3 processes a defective vertex, it yields either a rainbow coloring, or

187 a larger admissible coloring and the next defective vertex in a constant time. Finally, since  
 188 we start from an uncolored graph and color each additional vertex in a constant time, the  
 189 implied algorithm colors all vertices in  $O(n)$  time.  $\square$

190 For a partial coloring  $f$  of a graph  $G$  and a vertex  $v \in V(G)$ , we say that  $u \in V(G)$  is  
 191  $f$ -visible from  $v$ , if either  $vu \in E(G)$  or  $v$  and  $u$  have a common uncolored neighbor.

192 **Theorem 4.2.** *There exists a linear (in  $n$ ) algorithm finding an acyclic coloring for an*  
 193  *$n$ -vertex graph  $G$  using at most  $1 + \lfloor \frac{(1+\Delta(G))^3}{4} \rfloor$  colors.*

194 *Proof.* Let  $r := \Delta(G)$ . We start from the partial coloring  $f_0$  that has no colored vertices,  
 195 and for  $i = 1, \dots, n$  at Step  $i$  obtain a rainbow partial acyclic coloring  $f_i$  from  $f_{i-1}$  by  
 196 coloring one more vertex (without recoloring). The algorithm proceeds as follows: at Step  
 197  $i$  choose a vertex  $v_i$  with the most colored neighbors. Greedily color  $v_i$  with a color  $\alpha_i$  in  
 198  $C := \{1, \dots, 1 + \lfloor \frac{(1+r)^2}{4} \rfloor\}$  that is distinct from the colors of all vertices  $f_{i-1}$ -visible from  
 199  $v_i$ . We claim that we always can find such  $\alpha_i$  in  $C$ .

Suppose that at Step  $i$ ,  $v_i$  has exactly  $k$  colored neighbors. Then it has at most  $r - k$  un-  
 colored neighbors, and each of these uncolored neighbors has at most  $k$  colored neighbors.  
 So, the total number of vertices  $f_{i-1}$ -visible from  $v_i$  is at most

$$k + (r - k)k = k(r + 1 - k) \leq \lfloor \frac{(r + 1)^2}{4} \rfloor = |C| - 1,$$

200 and we can find a suitable color  $\alpha_i$  for  $v_i$ .

201 It now suffices to show that for each  $i$ , coloring  $f_i$  is rainbow and acyclic. For  $f_0$ , this  
 202 is obvious. Assume now that  $f_{i-1}$  is rainbow and acyclic. Since  $v_i$  is rainbow in  $f_{i-1}$ ,  
 203 coloring it with  $\alpha_i$  does not create bicolored cycles. Thus,  $f_i$  is acyclic. Also since  $\alpha_i$  is  
 204 distinct from the colors of all vertices  $f_{i-1}$ -visible from  $v_i$ ,  $f_i$  is rainbow.

205 For the runtime, note that at Step  $i$  the algorithm considers only  $v_i$  and vertices at  
 206 distance at most 2 from  $v_i$ . As in the proof of Theorem 4.1, it is sufficient to maintain  $r + 1$   
 207 databases each containing all vertices with a given number of colored neighbors. This  
 208 allows a constant time search for a vertex with the greatest number of colored neighbors.  
 209 Moving a vertex as its number of colored neighbors changes takes a constant amount of  
 210 time. Choosing and coloring  $v_i$  together with updating the databases then takes  $O(r^2)$   
 211 time. Hence the running time of the algorithm is at most  $c_r n$ , where  $c_r$  depends on  $r$ .  $\square$

## 212 5 Proof of Lemma 2.3

213 We will prove that under the conditions of the lemma, either its conclusion holds or there is  
 214 an admissible coloring  $f''$  larger than  $f$ . Since  $G$  is finite, repeating the argument eventually  
 215 yields either an acyclic coloring of the whole  $G$  or a rainbow coloring. In both cases we do  
 216 not have defective vertices.

217 Let  $H$  be the subgraph of  $G$  induced by the uncolored vertices. Let  $x$  be the sole  
 218 defective vertex under  $f$  and let  $y_1, y_2, \dots, y_5$  be its neighbors. By the definition of a  
 219 defective vertex,  $x$  has two neighbors of the same color. We will assume that  $f(y_1) =$   
 220  $f(y_2) = 1$  and that  $y_1$  is admissible. When more than two neighbors of  $x$  are colored, we  
 221 assume for  $i = 3, 4, 5$  that if  $y_i$  is colored, then  $f(y_i) = i - 1$ . Also for  $i = 1, \dots, 5$ , the  
 222 four neighbors of  $y_i$  distinct from  $x$  will be denoted by  $y_{i,1}, \dots, y_{i,4}$  (some vertices will  
 223 have more than one name, since they may be adjacent to more than one  $y_i$ ). We consider  
 224 several cases depending on  $d_H(x)$ .

225 **Case 1:**  $d_H(x) = 0$ . First we try to color  $x$  with colors 5, 6, and 7. If this is not  
 226 allowed, then for  $j = 5, 6, 7$ ,  $G$  has a 1,  $j$ -colored  $y_1, y_2$ -path. This forces that both of  $y_1$   
 227 and  $y_2$  have neighbors with colors 5, 6, and 7, each of which is adjacent to another vertex  
 228 of color 1. In particular, both  $y_1$  and  $y_2$  are admissible. For  $i = 1, 2$  and  $j = 1, 2, 3$ , we  
 229 suppose that  $f(y_{i,j}) = j + 4$  and  $y_{i,j}$  is adjacent to another vertex of color 1.

230 *Case 1.1:* For some  $i \in \{1, 2\}$ ,  $y_{i,4}$  is colored and  $f(y_{i,4}) \notin \{5, 6, 7\}$ . By symmetry,  
 231 we may assume that  $i = 1$  and  $f(y_{1,4}) = 2$ . Recolor  $y_1$  with 3 and call the new admissible  
 232 coloring  $f'$ . If we can now recolor  $y_2$  so that the resulting coloring  $f''$  is rainbow on  
 233  $G - xy_2 - xy_1$  or the only defective vertex in  $f''$  on  $G - xy_2 - xy_1$  is  $y_{2,4}$ , then we do  
 234 this recoloring and color  $x$  with 1. Since  $y_1$  and  $y_2$  have no neighbors of color 1 apart  
 235 from  $x$ , we obtained an admissible coloring of  $G$  larger than  $f$ . If we cannot recolor  $y_2$   
 236 to get such a coloring, then  $y_{2,4}$  is colored with a color  $c \in \{5, 6, 7\}$ . Moreover, in this  
 237 case by Lemma 2.1 applied to  $y_2$  in coloring  $f'$  of  $G - xy_2 - xy_1$ , we can change the  
 238 colors of only  $y_2$  and some  $y \in \{y_{2,1}, y_{2,2}, y_{2,3}, y_{2,4}\}$  to get an admissible coloring  $f_1$  of  
 239  $G - xy_2 - xy_1$ . Moreover, by Lemma 2.1,  $f_1(y) \in \{5, 6, 7\}$ . Then by coloring  $x$  with 1  
 240 we obtain a rainbow coloring of  $G$ , as above.

241 *Case 1.2:*  $y_{1,4}$  is not colored. By Lemma 2.2 for vertex  $y_1$  in  $G - xy_1$ , either  $G - xy_1$   
 242 has a rainbow coloring  $f'$  that differs from  $f$  only at  $y_1$  (in which case by symmetry, we  
 243 may assume that  $f'(y_1) = 3$  and proceed further exactly as in Case 1.1), or  $G - xy_1$  has a  
 244 larger rainbow coloring  $f'$  satisfying statements 1)–4) of Lemma 2.2. In particular, by 4),  
 245 none of  $y_2, y_3, y_4, y_5$  changes its color and  $y_1$  remains admissible. This finishes Case 1.2.

246 By the symmetry between  $y_1$  and  $y_2$ , the remaining subcase is the following.

247 *Case 1.3:*  $f(y_{1,4}) = 5$  and  $f(y_{2,4}) = c \in \{5, 6, 7\}$ . By Lemma 2.1 applied to  $y_1$   
 248 in  $G - xy_1$ , we can recolor  $y_1$  and at most one other vertex (a neighbor of  $y_1$ ) to obtain  
 249 another admissible coloring  $f'$ . If  $f'(y_1) \in \{5, 6, 7\}$ , then  $f'$  is a rainbow coloring, as  
 250 claimed. So, we may assume that  $f'(y_1) = c_1 \in \{2, 3, 4\}$ . If all the colors 5, 6, 7 are  
 251 present on neighbors of  $y_2$ , then again by Lemma 2.1 (applied now to  $y_2$  in coloring  $f'$  of  
 252  $G - xy_2$ ),  $G$  has an admissible coloring  $f''$  that differs from  $f'$  only at  $y_2$  and maybe at  
 253 one neighbor of  $y_2$ . Then coloring  $x$  with 1 we get a rainbow coloring. So, some color  
 254 in  $\{5, 6, 7\}$  is not present in  $f'(N(y_2))$ . By Lemma 2.1, this may happen only if  $y_{1,1}$  is a  
 255 common neighbor of  $y_1$  and  $y_2$ , and  $c = f(y_{2,4}) \neq 5$ . In particular, in this case,  $y_{1,1}$  has  
 256 neighbors of colors 1 (they are  $y_1$  and  $y_2$ ), 2, 3, and 4. Since  $c \neq 5$ , we may assume that  
 257  $c = 6$ . By the symmetry between  $y_1$  and  $y_2$ , we conclude that, in  $f$ , vertex  $y_{2,2}$  also is a  
 258 common neighbor of  $y_1$  and  $y_2$  and has neighbors of colors 1 (they are  $y_1$  and  $y_2$ ), 2, 3, and  
 259 4. Returning to coloring  $f'$ , we see that  $y_2$  has no neighbors of color 5, and its neighbors  
 260  $y_{1,1}$  (formerly of color 5) and  $y_{2,2}$  (by the previous sentence) also have no neighbors of  
 261 color 5. So, recoloring  $y_2$  with 5 yields an admissible coloring of  $G$ . Now coloring  $x$  with  
 262 1 creates a larger rainbow coloring.

263 **Case 2:**  $d_H(x) = 1$ . We first try to color  $x$  with 4. If no bicolored cycle is formed,  
 264 then either we have a rainbow coloring or an admissible coloring with defective vertex  
 265  $y_5$  and an admissible neighbor  $x$ . Hence we may assume that coloring  $x$  with 4 creates  
 266 a bicolored cycle. This then gives each of  $y_1$  and  $y_2$  a neighbor of color 4. A similar  
 267 argument gives each of  $y_1$  and  $y_2$  a neighbor of color 5, 6, and 7, i.e., both  $y_1$  and  $y_2$   
 268 are rainbow. Recoloring  $y_1$  with color 2 allows us to repeat the argument at  $y_3$ . Then  
 269  $y_3$  also has neighbors of each of the colors 4, 5, 6, and 7. If  $y_5$  has no neighbor of color  
 270 2, then recoloring (in the original coloring  $f$ )  $y_3$  with 1, and coloring  $x$  with 2 yields a  
 271 rainbow coloring. So, by the symmetry between colors 1, 2, and 3, we may assume that for

272  $i \in \{1, 2, 3\}$ ,  $f(y_{5,i}) = i$ . Since  $y_5$  is rainbow, by the symmetry between colors 4, 5, 6,  
 273 and 7, we may assume that either  $f(y_{5,4}) = 4$ , or  $y_{5,4}$  is not colored. In both cases, recolor  
 274 (in the original coloring  $f$ )  $y_3$  with 1, color  $x$  with 2 and  $y_5$  with 5. We get an admissible  
 275 coloring larger than  $f$ , where only  $y_{5,4}$  may be defective.

276 **Case 3:**  $d_H(x) = 3$ . If one of the uncolored neighbors  $y_3, y_4, y_5$  (say,  $y_3$ ) of  $x$  has 4  
 277 colored neighbors, then we may color  $y_3$  with some  $c \notin f(N(y_3)) \cup \{1\}$  and thus create  
 278 an admissible coloring larger than  $f$ . Hence we may assume that each of  $y_3, y_4$ , and  $y_5$  has  
 279 at most 3 colored neighbors.

280 *Case 3.1:* One of  $y_1$  and  $y_2$  has three neighbors of different colors such that each of  
 281 these neighbors has another neighbor of color 1. Suppose for example that for  $j = 1, 2, 3$ ,  
 282  $f(y_{1,j}) = 1 + j$  and  $y_{1,j}$  has another neighbor of color 1. If  $y_1$  has a fourth color, say  $c$ ,  
 283 in its neighborhood, then we recolor  $y_1$  with a color  $c' \notin \{1, c, 5, 6, 7\}$  and get a rainbow  
 284 coloring of  $G$ . Suppose now that color  $c \in \{5, 6, 7\}$  appears twice on  $N(y_1)$ . Then by  
 285 Lemma 2.1 applied to  $y_1$  in  $G - xy_1$ , we can change the color of  $y_1$  and at most one other  
 286 vertex that is a neighbor of  $y_1$  not adjacent to uncolored vertices to get another rainbow  
 287 coloring of  $G - xy_1$ . Then this coloring will also be a rainbow coloring of  $G$ . Finally,  
 288 suppose that  $y_1$  has an uncolored neighbor  $y_{1,4}$ . Applying Lemma 2.2 to  $y_1$  in  $G - xy_1$   
 289 we either recolor only  $y_1$  and get a rainbow coloring of  $G$  (finishing the case), or obtain  
 290 a rainbow coloring  $f'$  of  $G - xy_1$  larger than  $f$  satisfying the conclusions of the lemma.  
 291 Since each of  $y_3, y_4$  and  $y_5$  has at least two neighbors left uncolored by  $f$ , none of them  
 292 may play role of  $z_4$  or  $z_{4,4}$  in Lemma 2.2 when they get colored. Then  $f'$  is an admissible  
 293 coloring of  $G$  where only  $x$  could be a defective vertex with admissible neighbor  $v$ . This  
 294 proves Case 3.1.

295 Let  $T$  be the set of colors  $c$  such that more than one of the vertices  $y_3, y_4$  and  $y_5$  has a  
 296 neighbor of color  $c$ . Since  $y_3, y_4$  and  $y_5$  have in total at most 9 colored neighbors,  $|T| \leq 4$ .

297 *Case 3.2:*  $|T| \leq 3$ . By symmetry, we may assume that  $T \subseteq \{2, 3, 4\}$ . If coloring  $x$   
 298 with  $c \in \{5, 6, 7\}$  does not create a bicolored cycle, then it will yield an admissible coloring  
 299 larger than  $f$ . So, we may assume that each of  $y_1$  and  $y_2$  has in its neighborhood vertices  
 300 of colors 5, 6, and 7, each of which is adjacent to another vertex of color 1. So, we have  
 301 Case 3.1.

302 *Case 3.3:*  $|T| = 4$ . Let  $T = \{2, 3, 4, 5\}$ . As in Case 3.1, we may assume that each of  
 303  $y_1$  and  $y_2$  is adjacent to vertices of colors 6 and 7, each of which have another neighbor of  
 304 color 1.

305 Let  $y_3$  have exactly 3 colored neighbors labeled  $y_{3,1}, y_{3,2}, y_{3,3}$  with colors 2, 3, 4. Let  
 306  $y_{3,4}$  be the uncolored neighbor of  $y_3$ . Then if  $y_{3,4}$  has no neighbor of color 5, we may color  
 307  $y_3$  with 5 to get a new admissible coloring. Hence  $y_{3,4}$  is adjacent to a vertex of color 5.  
 308 Similarly,  $y_{3,4}$  has neighbors of color 6 and 7. By symmetry, we may assume that a vertex  
 309 of color 2 is adjacent to at most one of  $y_4$  and  $y_5$ .

310 *Case 3.3.1:*  $y_{3,4}$  has no neighbor of color 1. We try to color  $y_3$  with 1 and  $x$  with 2.  
 311 If this does not produce a new admissible coloring, then one of  $y_1$  or  $y_2$ , say  $y_1$ , has a  
 312 neighbor of color 2 that is adjacent to another vertex of color 1. So, we again get Case 3.1.

313 *Case 3.3.2:*  $y_{3,4}$  has a neighbor of color 1. If  $y_{3,1}$  has no neighbor of color 1, then we  
 314 again try to color  $y_3$  with 1 and  $x$  with 2, but also color  $y_{3,4}$  with 2. Then we simply repeat  
 315 the argument of Case 3.3.1. So, suppose that  $y_{3,1}$  has a neighbor of color 1. If  $y_{3,1}$  has no  
 316 neighbor of some color  $\alpha \in \{5, 6, 7\}$ , then we color  $y_{3,4}$  with 2 and  $y_3$  with  $\alpha$ . Thus  $y_{3,1}$   
 317 has neighbors of colors 1, 5, 6, 7. Then we recolor  $y_{3,1}$  with 3 and color  $y_3$  with 2.






Figure 1: Cases 4.3 and 4.3.4.3 from the proof of Lemma 2.3.

318 **Case 4:**  $d_H(x) = 2$ . As at the beginning of Case 3, we conclude that each of the  
 319 uncolored vertices  $y_4$  and  $y_5$  has at least one uncolored neighbor besides  $x$ .

320 Let  $B$  be the set of colors appearing in the neighborhoods of both,  $y_4$  and  $y_5$ . By the  
 321 previous paragraph,  $|B| \leq 3$ .

322 *Case 4.1:*  $|B| \leq 1$ . We may assume that  $\{4, 5, 6, 7\} \cap B = \emptyset$ . Try to color  $x$  with  
 323 4. By the definition of  $B$ , either a two-colored cycle appears, or we get a new admissible  
 324 coloring larger than  $f$ . Hence we may assume that coloring  $x$  with 4 creates a bicolored  
 325 cycle. Since this cycle necessarily goes through  $y_1$ ,  $y_1$  is adjacent to a vertex with color  
 326 4. Similarly,  $y_1$  is adjacent to vertices with colors 5, 6, and 7. Then recoloring  $y_1$  with 3  
 327 yields a rainbow coloring of  $G$ .

328 *Case 4.2:*  $|B| = 2$ . If  $1 \in B$  or  $2 \in B$ , then the argument of Case 4.1 holds. Assume  
 329 that  $B = \{3, 4\}$ . Similarly to Case 4.1, we may assume that for  $i = 1, 2$  and  $j = 1, 2, 3$ ,  
 330  $y_i$  is adjacent to a vertex  $y_{i,j}$  of color  $j + 4$  that is adjacent to another vertex of color 1 (in  
 331 particular,  $y_1$  and  $y_2$  may have a common neighbor of color  $j + 4$ ).

332 If  $y_1$  is rainbow, then uncoloring  $y_1$  and coloring  $x$  with 7 gives Case 1 or Case 2. Thus  
 333 we may assume that  $y_1$  and (by symmetry)  $y_2$  are not rainbow. So, we may assume that  
 334 for  $i = 1, 2$ , the fourth neighbor  $y_{i,4}$  of  $y_i$  distinct from  $x$  has color  $c_i \in \{5, 6, 7\}$ . By  
 335 symmetry, we may assume that  $c_1 = 5$ . Similarly to Case 1.3, by Lemma 2.1 applied to  
 336  $y_1$  in  $G - xy_1$ , we can recolor  $y_1$  and at most one other vertex (a neighbor of  $y_1$ ) to obtain  
 337 another rainbow coloring  $f'$  of  $G - xy_1$ . If  $f'(y_1) \in \{3, 4, 5, 6, 7\}$ , then  $f'$  is a rainbow  
 338 coloring of  $G$ , as claimed. So, we may assume that  $f'(y_1) = 2$ . Now practically repeating  
 339 the argument of Case 1.3, we find a promised coloring.

340 *Case 4.3:*  $|B| = 3$  (see Figure 1). If  $2 \in B$ , then we can repeat the argument of Case  
 341 4.2 for  $B' = B - \{2\}$ . Hence we may assume that  $B \subseteq \{1, 3, 4, 5, 6, 7\}$ .

342 *Case 4.3.0:*  $1 \in B$ . Let  $B = \{1, 3, 4\}$ . Then some color in  $\{5, 6, 7\}$ , say 7, is not  
 343 present on  $N(y_4) \cup N(y_5)$ . Again, we may assume that for  $i = 1, 2$  and  $j = 1, 2, 3$ ,  $y_i$  is  
 344 adjacent to a vertex  $y_{i,j}$  of color  $j + 4$  that is adjacent to another vertex of color 1. If  $y_1$  is  
 345 rainbow, then we may uncolor  $y_1$  and color  $x$  with 7 to get Case 1 or Case 2. Suppose now

346 that  $y_1$  and  $y_2$  are not rainbow. By Lemma 2.1 applied to  $y_1$  in  $G - xy_1$ , we can recolor  $y_1$   
 347 and at most one other vertex (a neighbor of  $y_1$ ) to obtain another admissible coloring  $f'$ .  
 348 If  $f'(y_1) \in \{3, 4, 5, 6, 7\}$ , then  $f'$  is a rainbow coloring, as claimed. So, we may assume  
 349 that  $f'(y_1) = 2$ . But then we can use the argument of Case 4.2 with the roles of  $y_3$  and  $y_2$   
 350 switched. This proves Case 4.3.0.

351 So, from now on,  $B = \{3, 4, 5\}$ . For  $i = 4, 5$  and  $j = 1, 2, 3$ , let  $y_{i,j}$  be the neighbor of  
 352  $y_i$  of color  $j + 2$ . We write *the* neighbor, since  $y_4$  and  $y_5$  are rainbow. As observed at the  
 353 beginning of Case 4,  $y_4$  and  $y_5$  each have another uncolored neighbor, call them  $y_{4,4}$  and  
 354  $y_{5,4}$ . In particular,  $y_4$  and  $y_5$  have no neighbors colored with 6 or 7. If  $x$  can be colored with  
 355 either of 6 or 7 without creating a two-colored cycle, then we obtain a rainbow coloring.  
 356 Hence we assume that for  $i = 1, 2$  and  $j = 1, 2$ ,  $f(y_{i,j}) = j + 5$  and  $y_{i,j}$  has a neighbor of  
 357 color 1 distinct from  $y_i$ .

358 *Case 4.3.1:* One of  $y_1$  or  $y_2$ , say  $y_1$ , is rainbow. If  $y_{4,4}$  has no neighbor of color  
 359  $c \in \{6, 7\}$ , then we can color  $y_4$  with  $c$ , a contradiction to the maximality of  $f$ . If  $y_{4,4}$  has  
 360 no neighbor of color  $c' \in \{1, 2\}$ , then by uncoloring  $y_1$  and coloring  $y_4$  with  $c'$  and  $x$  with  
 361 6, we obtain an admissible coloring larger than  $f$ . So,  $f(N(y_{4,4})) = \{1, 2, 6, 7\}$ . Then we  
 362 may color  $y_{4,4}$  with 3 and uncolor  $y_1$  to get a new admissible coloring as large as  $f$  with  
 363 one defective vertex  $y_4$ , for which Case 2 holds. This finishes Case 4.3.1.

364 So, below  $y_1$  and  $y_2$  are not rainbow and hence each of them is adjacent to at least three  
 365 colored vertices.

366 *Case 4.3.2:* One of  $y_1$  or  $y_2$ , say  $y_1$ , is adjacent to an uncolored vertex  $y_{1,4} \neq x$ . We  
 367 may assume that  $f(y_{1,1}) = f(y_{1,2}) = 6$  and  $f(y_{1,3}) = 7$ . First, we try to color  $x$  with 7  
 368 and  $y_1$  with 3. Since the new coloring has at most one defective vertex, we may assume  
 369 that a two-colored cycle is created. Hence each of  $y_{1,1}$  and  $y_{1,2}$  is adjacent to a vertex of  
 370 color 3. The same argument gives these vertices neighbors of colors 4 and 5. Recall that  
 371 one of  $y_{1,1}$  and  $y_{1,2}$ , say  $y_{1,1}$ , has another neighbor of color 1. Then recoloring  $y_{1,1}$  with 2  
 372 gives an admissible coloring in which  $y_1$  is rainbow. Hence Case 4.3.1 applies to this new  
 373 coloring.

374 So, from now on each of  $y_1$  and  $y_2$  has 4 colored neighbors. Since  $y_1$  is admissible we  
 375 may assume w.l.o.g. that  $y_1$  is adjacent either to the colors 5, 6, 6, 7 or the colors 5, 5, 6, 7.

376 *Case 4.3.3:*  $y_1$  has one neighbor of color 5 and three neighbors with colors 6 or 7. We  
 377 may assume that  $f(y_{1,1}) = 5$ ,  $f(y_{1,2}) = f(y_{1,3}) = 6$ , and  $f(y_{1,4}) = 7$ . If coloring  $y_1$   
 378 with 3 or 4 yields an admissible coloring, then we are done; so we may assume that a two-  
 379 colored cycle is formed in each case. It follows that each of  $y_{1,2}$  and  $y_{1,3}$  has neighbors  
 380 colored with 3 and 4. By the symmetry between  $y_{1,2}$  and  $y_{1,3}$ , we may assume that  $y_{1,3}$   
 381 has a neighbor of color 1 other than  $y_1$ . If  $y_{1,3}$  is almost rainbow, then we can uncolor it,  
 382 recolor  $y_1$  with 3, and color  $x$  with 7: this will give an admissible coloring with the same  
 383 number of colored vertices as in  $f$ , and the only defective vertex  $y_{1,3}$ . Then either Case 1  
 384 or Case 2 applies to this new coloring. Hence we may assume that  $y_{1,3}$  has two neighbors  
 385 other than  $y_1$  that receive the same color. Then since  $y_{1,3}$  has no neighbor of color 2,  $y_1$   
 386 may now be recolored with color 2 without creating a bicolored cycle. Repeating the above  
 387 argument we derive that  $y_{1,2}$  has neighbors of colors 2, 3, and 4, and one of these colors  
 388 appears twice on  $N(y_{1,2}) - y_1$ . By Lemma 2.1 applied to  $y_{1,3}$  in the graph  $G - y_{1,3}y_1$  for  
 389 the original coloring, we can change its color and the color of at most one other vertex (that  
 390 is a neighbor of  $y_{1,3}$ , all of whose neighbors are colored) to get an admissible coloring of  
 391  $G - y_{1,3}y_1$ . Since  $y_2$  and  $y_3$  are adjacent to the uncolored vertex  $x$ , their colors are not  
 392 changed. If  $y_{1,3}$  receives color 1, then we recolor  $y_1$  with 3 and get a rainbow coloring of

393  $G$ . If  $y_{1,3}$  receives a color other than 1, then we color  $x$  with 6 and again get a rainbow  
394 coloring of  $G$ .

395 *Case 4.3.4:*  $y_1$  has two neighbors of color 5 (see Figure 1). We may assume that  
396  $f(y_{1,1}) = f(y_{1,2}) = 5$ ,  $f(y_{1,3}) = 6$ , and  $f(y_{1,4}) = 7$ . If  $y_1$  can be recolored with either  
397 3 or 4, this would give a rainbow coloring  $f'$ . Hence we assume that both of  $y_{1,1}$  and  $y_{1,2}$   
398 are adjacent to vertices with colors 3 and 4.

399 *Case 4.3.4.1:* One of  $y_{1,1}$  or  $y_{1,2}$ , say  $y_{1,1}$ , is rainbow. Then uncoloring  $y_{1,1}$  and  
400 coloring  $y_1$  with 3 and  $x$  with 7 yields either a rainbow coloring  $f'$  or a new admissible  
401 coloring (with the same number of colored vertices) with the defective vertex  $y_{1,1}$  and  
402 admissible colored neighbor  $y_1$ . In the former case, we are done. In the latter, if one of the  
403 previous cases occurs, then we are done again. So, we may assume that Case 4.3.4 occurs.  
404 By the symmetry between colors 3 and 4, we may assume that apart from  $y_1$ , vertex  $y_{1,1}$   
405 has a neighbor of color 3, a neighbor of color 4, and two uncolored neighbors, say  $z_1$  and  
406  $z_2$ , each of whose has another uncolored neighbor and 3 colored neighbors. Moreover, the  
407 same 3 colors appear on the neighborhoods of  $z_1$  and  $z_2$ , and since Case 4.3.4 holds, by  
408 the symmetry between colors 6 and 7, both of them are among these 3 colors. Then either  
409 coloring  $y_{1,1}$  with 1 yields a rainbow coloring or coloring  $y_{1,1}$  with 2 does.

410 *Case 4.3.4.2:* Each of  $y_{1,1}$  and  $y_{1,2}$  has a neighbor of color 2 that has another neighbor  
411 of color 5. Since  $y_{1,1}$  is not rainbow, the fourth neighbor of  $y_{1,1}$  has color  $c \in \{2, 3, 4\}$ .  
412 Since  $y_1$  cannot be recolored with 3 or 4, some neighbor, say  $r$ , of  $y_{1,1}$  of color  $c$  has  
413 another neighbor of color 5. If in the graph  $G - y_1 y_{1,1}, y_{1,1}$  can be recolored with 1, then  
414 we may recolor  $y_1$  with 3 and get a rainbow coloring of  $G$ . If  $y_{1,1}$  can be recolored with  
415 either of 6 or 7, then we have Case 4.3.3. To disallow coloring  $y_{1,1}$  with 1, 6, and 7,  $r$  must  
416 be adjacent to vertices with each of these colors. By the symmetry between colors 3 and  
417 4, we assume that  $f(r) \neq 4$ . If the neighbor  $r'$  of  $y_{1,1}$  with  $f(r') = 4$  has no neighbor of  
418 color  $c' \in \{6, 7\}$ , then we recolor  $r$  with 4 and  $y_{1,1}$  with  $c'$  thus getting Case 4.3.3. If  $r'$   
419 has no neighbor of color 1, then we recolor  $r$  with 4,  $y_{1,1}$  with 1, and  $y_1$  with 3 obtaining a  
420 rainbow coloring. Finally if  $f(N(r') - y_{1,1}) = \{1, 5, 6, 7\}$ , then we recolor  $r'$  with 3,  $y_{1,1}$   
421 with 4, and  $y_1$  with 3.

422 The last subcase is:

423 *Case 4.3.4.3:*  $y_{1,1}$  has no neighbor of color 2 that has another neighbor of color 5.  
424 Then recoloring  $y_1$  with 2 creates another admissible coloring  $f'$ . We may then repeat our  
425 previous argument with  $y_3$  playing the role of  $y_2$  to conclude that  $y_3$  has neighbors of color  
426 6 and 7. If  $y_3$  is admissible, then repeating the above argument we conclude that  $y_3$  may be  
427 recolored with color 1 in the original coloring  $f$ . Then after this recoloring, by coloring  $x$   
428 with 2 we get a rainbow coloring. Also, if  $y_2$  is admissible in  $f$ , then we may recolor both  
429 of  $y_1$  and  $y_2$  with 2 and color  $x$  with 1 to get a rainbow coloring. Hence we may assume  
430 that all the neighbors of  $y_2$  and  $y_3$  apart from  $x$  are colored with 6 or 7. Recall that for  
431  $i = 4, 5$  and  $j = 1, 2, 3$ ,  $f(y_{i,j}) = j + 2$  and  $y_{i,4}$  is uncolored. If for some  $i \in \{4, 5\}$ ,  $y_{i,4}$   
432 has no neighbor of color  $c \in \{6, 7\}$ , then we can color  $y_i$  with  $c$  and get a better admissible  
433 coloring. Since none of  $y_1, y_2$ , or  $y_3$  has a neighbor with color 3, if  $y_{4,4}$  has no neighbor of  
434 color 1 or  $y_{5,4}$  has no neighbor of color 2, then by coloring  $y_4$  with 1,  $y_5$  with 2 and  $x$  with  
435 3 creates an admissible coloring with more colored vertices. By the symmetry between  
436 colors 1 and 2, each of  $y_{4,4}$  and  $y_{5,4}$  has neighbors of colors 1, 2, 6, and 7.

437 If  $y_{4,1}$  does not have a neighbor of color  $c' \in \{1, 2, 6, 7\}$ , then coloring  $y_{4,4}$  with 3,  $y_4$   
438 with  $c'$  and  $x$  with 4 yields an admissible coloring. Otherwise, we recolor  $y_{4,1}$  with 4 and  
439 color  $y_4$  with 3. This proves the lemma.

## 6 Acknowledgment

We thank the referees for the helpful comments.

## References

- [1] M. O. Albertson and D. M. Berman, The acyclic chromatic number, *Congressus Numerantium*, No. XVII (1976), Utilitas Math., Winnipeg, Man., 51–69. 153, 154
- [2] M. O. Albertson and D. M. Berman, Every planar graph has an acyclic 7-coloring, *Israel J. Math.* **28** (1977), 169–174. 153
- [3] M. O. Albertson and D. M. Berman, An acyclic analogue to Heawood’s theorem, *Glasgow Math. J.* **19** (1978), 163–166. 153
- [4] M. O. Albertson, G. G. Chappell, H. A. Kierstead, A. Kündgen and R. Ramamurthi, Coloring with no 2-colored  $P_4$ ’s, *Electron. J. Combin.* **11** (2004), R#26. 153
- [5] N. Alon, C. McDiarmid and B. Reed, Acyclic coloring of graphs, *Random Structures Algorithms* **2** (1991), 277–288. 154
- [6] M. I. Burstein, Every 4-valent graph has an acyclic 5-coloring, *Sobšč. Akad. Nauk Gruzin. SSR* **93** (1979), 21–24. 154
- [7] G. Fertin and A. Raspaud, Acyclic coloring of graphs of maximum degree five: nine colors are enough, *Information Processing Letters* **105** (2008), 65–72. 154
- [8] B. Grünbaum, Acyclic colorings of planar graphs, *Israel J. Math.* **14** (1973), 390–408. 153
- [9] H. Hocquard, M. Montassier, Acyclic coloring of graphs with maximum degree five, <http://hal.archives-ouvertes.fr/hal-00375166/en/>. 154
- [10] A. V. Kostochka, *Upper bounds on the chromatic characteristics of graphs*, PhD Thesis, Institute of Mathematics, Novosibirsk, 1978. 154
- [11] K. Kothapalli, V. Satish and V. Ch. Venkaiah, Acyclic vertex coloring of graphs of maximum degree  $\Delta$ , Talk at 75th Annual conference of the Indian Math. Society, Kalasalingam Univ., Krishnakoil, December 2009. 154
- [12] K. Kothapalli, S. Varagani, V. Ch. Venkaiah and K. Yadav, Acyclic Vertex Coloring of Graphs of Maximum Degree 5, Proc. Int. Conf. on Graph Theory and its Applications, Amrita Vishwa Vidyapeetham, Coimbatore, December, 2008. 154